

LAX OPERATOR ALGEBRAS

IGOR M. KRICHEVER AND OLEG K. SHEINMAN

ABSTRACT. In this paper we develop a general concept of Lax operators on algebraic curves introduced in [1]. We observe that the space of Lax operators is closed with respect to their usual multiplication as matrix-valued functions. We construct the orthogonal and symplectic analogs of Lax operators, prove that they constitute almost graded Lie algebras and construct local central extensions of those Lie algebras.

1. INTRODUCTION

A general concept of Lax operators on algebraic curves was proposed by one of the authors in [1], where the theory of conventional Lax and zero curvature representations with the *rational spectral parameter* was generalized on the case of algebraic curves Γ of arbitrary genus g . The linear space of such operators associated with an effective divisor $D = \sum_k n_k P_k$, $P_k \in \Gamma$ was defined as the space of meromorphic $(n \times n)$ matrix-valued functions on Γ having poles of multiplicity at most n_k at the points P_k , and at most simple poles at ng points γ_s more. The coefficients of the Laurent expansion of those matrix-valued functions in the neighborhood of a point γ_s had to obey certain linear constraints parametrized by the point α_s of a certain projective space (see relations (2.1)-(2.3) below).

According to [12], the generic sets (γ_s, α_s) parameterize stable rank n degree ng framed holomorphic vector bundles B on Γ . In [1] it was noticed that the requirements on the form of Lax operators at the points γ_s mean that these operators can be seen as meromorphic sections of the bundle $End(B)$ with D as the divisor of poles. As an easy consequence of this remark, we have that Lax operators having poles of arbitrary orders at the points P_k form an algebra with respect to the usual pointwise multiplication.

In the simplest case of two marked points, $D = P_+ + P_-$, this enables us to define in the algebra of the corresponding operators an almost graded structure generalizing the graded structure of the classic affine algebra $\widehat{\mathfrak{gl}(n)}$. Recall that a Lie algebra $\mathcal{V}$ is called *almost graded* if $\mathcal{V} = \oplus \mathcal{V}_i$ where

Research is supported in part by National Science Foundation under the grant DMS-04-05519 (I.M.Krichever), by Russian Foundation for Basic Research under the grant 05-01-00170, and by the program "Mathematical Methods in Nonlinear Dynamics" of the Russian Academy of Science (O.K.Sheinman).

$\dim \mathcal{V}_i < \infty$ and $[\mathcal{V}_i, \mathcal{V}_j] \subseteq \bigoplus_{k=i+j-k_0}^{k=i+j+k_1} \mathcal{V}_k$ where k_0 and k_1 do not depend on i, j .

A general notion of almost graded algebras and modules over them was introduced in [3]-[5] where the generalizations of Heisenberg and Virasoro algebras were introduced. In the series of works, see [11] for the review, the almost graded analogues of classic affine Lie algebras, called Krichever-Novikov (KN) current algebras, were investigated. It is natural to consider the algebra of Lax operators having poles at two points as a generalization of the KN $\mathfrak{gl}(n)$ -algebra.

A central extension of $\mathcal{V}$ is called *local* if it is an almost graded Lie algebra itself. Local central extensions are given by local 2-cocycles. A 2-cocycle γ is called local if a $K \in \mathbb{Z}$ exists such that $\gamma(\mathcal{V}_i, \mathcal{V}_j) = 0$ for $|i+j| > K$. The notion of local cocycle is introduced in [3]. In the same place its uniqueness was conjectured and the proof for the Virasoro-type algebras was outlined; a complete proof is given in [9, 10]. The locality condition is important in consideration of analogues of highest-weight representations.

To construct the orthogonal and symplectic analogues of Lax operators is the main goal of the present paper. In these cases, the Lax operators do not form any associative algebra, they only form a Lie algebra. For all classic Lie algebras $\mathfrak{g}$ the corresponding Lax operator algebras can be considered as a "twisted" version of the Krichever-Novikov current algebras and loop algebras.

In Section 2 we prove the multiplicative properties of $\mathfrak{gl}(n)$ -valued Lax operators, introduce $\mathfrak{g}$ -valued Lax operators for $\mathfrak{g} = \mathfrak{so}(n)$ and $\mathfrak{g} = \mathfrak{sp}(2n)$ and prove that they are closed with respect to the pointwise bracket.

In Section 3, we establish the almost graded structure on Lax operator algebras, and show that $\dim \mathcal{V}_i = \dim \mathfrak{g}$, as for KN algebras.

In Section 4, on every type of a Lax operator algebra, we define a 2-cocycle and prove its locality.

Authors are thankful to M.Schlichenmaier for his fruitful criticism.

2. LAX OPERATORS AND THEIR LIE BRACKET

2.1. Lax operator algebras for $\mathfrak{gl}(n)$ and $\mathfrak{sl}(n)$. Following [1], call by a *Lax operator* with the Tyurin parameters $\{\alpha_s, \gamma_s | s = 1, \dots, gr\}$ a $\mathfrak{gl}(n)$ -valued function L on Γ , holomorphic outside $P_{\pm}$ and $\{\gamma_s | s = 1, \dots, gr\}$, having at the latters at most simple poles:

$$(2.1) \quad L = \frac{L_{s,-1}}{z - z_s} + L_{s0} + O(z - z_s), \quad z_s = z(\gamma_s),$$

and such that

(i) $L_{s,-1} = \alpha_s \beta_s^t$ where β_s^t is a row, and

$$(2.2) \quad \text{tr } L_{s,-1} = \beta_s^t \alpha_s = 0,$$

where $\alpha_s \in \mathbb{C}^n$ is fixed, $\beta_s \in \mathbb{C}^n$ is arbitrary, and the upper t denotes a transposed matrix. In particular, $L_{s,-1}$ has rank 1;

(ii) α_s is an eigenvector of the matrix L_{s0} :

$$(2.3) \quad L_{s0} \alpha_s = k_s \alpha_s.$$

Lemma 2.1. *Let L' and L'' satisfy the conditions (2.1)-(2.3). Then $L = L' L''$ also satisfies them.*

Proof. From (2.1) we have

$$(2.4) \quad L = \frac{L'_{s,-1} L''_{s,-1}}{(z - z_s)^2} + \frac{L'_{s,-1} L''_{s0} + L'_{s0} L''_{s,-1}}{(z - z_s)} + L'_{s,-1} L''_{s1} + L'_{s0} L''_{s0} + L'_{s1} L''_{s,-1} + O(1).$$

From (2.2) for L' it follows that the first term vanishes:

$$L'_{s,-1} L''_{s,-1} = \alpha_s (\beta_s'^t \alpha_s) \beta_s''^t = 0.$$

For the second one we have $L_{s,-1} = L'_{s,-1} L''_{s0} + L'_{s0} L''_{s,-1} = \alpha_s (\beta_s'^t L''_{s0}) + (L'_{s0} \alpha_s) \beta_s''^t$. By (2.2) for L' it follows $L'_{s0} \alpha_s = k'_s \alpha_s$, hence $L_{s,-1} = \alpha_s \beta_s^t$ where $\beta_s^t = \beta_s'^t L''_{s0} + k'_s \beta_s''^t$. Further on, $\text{tr } L_{s,-1} = (\beta_s'^t L''_{s0} + k'_s \beta_s''^t) \alpha_s = k_s'' \beta_s'^t \alpha_s + k'_s \beta_s''^t \alpha_s = 0$.

Consider the expression $L_{s,0} \alpha_s$, where $L_{s,0} = L'_{s,-1} L''_{s1} + L'_{s0} L''_{s0} + L'_{s1} L''_{s,-1}$. From the definition of the Lax operators it follows that $L''_{s,-1} \alpha_s = 0$ and $L'_{s0} L''_{s0} \alpha_s = k'_s k_s'' \alpha_s$. We also have $L'_{s,-1} L''_{s1} \alpha_s = \alpha_s (\beta_s'^t L''_{s1} \alpha_s)$. Hence α_s is an eigenvector of the matrix $L_{s,0}$ with the eigenvalue $k_s = \beta_s'^t L''_{s1} \alpha_s + k'_s k_s''$. $\square$

Since the conditions (2.1),(2.2) are linear, the Lax operators constitute an associative algebra, hence the corresponding Lie algebra. The latter is called *Lax operator algebra*.

If the function L , besides (2.1)–(2.3), satisfies the condition $\text{tr } L = 0$, it is called an $\mathfrak{sl}(n)$ -valued Lax operator. Such Lax operators form a Lie algebra.

2.2. Lax operator algebras for $\mathfrak{so}(n)$. For the elements of this Lie algebra we have $X^t = -X$. We introduce the matrix-function L taking values in $\mathfrak{so}(n)$ by the same expansion as in the Section 2.1 but change the requirement (i) of that section for the reason that there is no rank one skew-symmetric matrix, and change accordingly the requirement (ii). We

omit the index s for brevity and write down the expansion (2.1) in the form

$$(2.5) \quad L = \frac{L_{-1}}{z} + L_0 + O(z)$$

where $L_0, L_1, \dots$ are skew-symmetric. Instead the condition (i) of Section 2.1 we require

$$(2.6) \quad L_{-1} = \alpha\beta^t - \beta\alpha^t$$

where $\alpha \in \mathbb{C}^n$ is fixed, $\beta \in \mathbb{C}^n$ is arbitrary and

$$(2.7) \quad \alpha^t\alpha = \beta^t\alpha (= \alpha^t\beta) = 0.$$

Likewise (2.3) we require

$$(2.8) \quad L_0\alpha = k\alpha$$

for some complex number k .

We will prove now the closeness of the space of Lax operators with respect to the Lie bracket in case of $\mathfrak{g} = \mathfrak{so}(n)$ in question. We stress that there is no structure of associative algebra in this case.

Lemma 2.2. *The properties (2.5)–(2.8) are invariant with respect to the Lie bracket.*

Proof. 1). First, prove the absence of the term with z^{-2} . The corresponding coefficient is equal to

$$\begin{aligned} [L_{-1}, L'_{-1}] &= [\alpha\beta^t - \beta\alpha^t, \alpha\beta'^t - \beta'\alpha^t] \\ &= (\beta^t\alpha)(\alpha\beta'^t - \beta'\alpha^t) - (\alpha^t\alpha)(\beta'\beta^t - \beta\beta'^t) - (\alpha^t\beta')(\alpha\beta^t - \beta\alpha^t). \end{aligned}$$

It vanishes due to (2.7) (applied to both β and β'). We stress that for the product $L_{-1}L'_{-1}$ the term with z^{-2} does not vanish.

2). Now calculate the term with z^{-1} in the product LL' . The coefficient is equal to

$$\begin{aligned} L_{-1}L'_0 + L_0L'_{-1} &= \alpha(\beta^tL'_0) - \beta(\alpha^tL'_0) + (L_0\alpha)\beta'^t - (L_0\beta')\alpha^t \\ &= \alpha(\beta^tL'_0) - \beta(-k'\alpha^t) + k\alpha\beta'^t - (L_0\beta')\alpha^t \\ &= \alpha(\beta^tL'_0 + k\beta'^t) - (L_0\beta' - k'\beta)\alpha^t \end{aligned}$$

(here we used the relation (2.8)). We see that it is not of the required form (2.6). Now consider the corresponding term of the commutator of L and L' . Then we obtain

$$\begin{aligned} [L, L']_0 &= \alpha(\beta^tL'_0 - \beta'^tL_0 + k\beta'^t - k'\beta^t) - (L_0\beta' - L'_0\beta - k'\beta + k\beta')\alpha^t \\ &= \alpha\beta'^t - \beta''\alpha^t \end{aligned}$$

where $\beta''' = \beta^t L'_0 - \beta'^t L_0 + k\beta'' - k'\beta^t$.

It is easy to check that β'' satisfies (2.7).

3) Check the requirement (2.8) on the eigenvalue of the degree 0 matrix coefficient $(LL')_0 = L_{-1}L'_1 + L_0L'_0 + L_1L'_{-1}$.

By (2.7) $L'_{-1}\alpha = 0$, hence for the third term we have $L_1L'_{-1}\alpha = 0$.

For the first term we have

$$L_{-1}L'_1\alpha = (\alpha\beta^t - \beta\alpha^t)L'_1\alpha = \alpha(\beta^t L'_1\alpha) - \beta(\alpha^t L'_1\alpha).$$

The last term on the right hand side of this equation vanishes due to skew-symmetry of L'_1 .

Thus

$$(2.9) \quad (LL')_0\alpha = k''\alpha \quad \text{where} \quad k'' = \beta^t L'_1\alpha + kk'.$$

□

2.3. Lax operator algebras for $\mathfrak{sp}(2n)$. For the elements of the symplectic algebra we have $X^t = -\sigma X \sigma^{-1}$ where σ is a nondegenerate skew-symmetric matrix.

Take the expansion for L in the form

$$(2.10) \quad L = \frac{L_{-2}}{z^2} + \frac{L_{-1}}{z} + L_0 + L_1z + L_2z^2 + O(z^3)$$

(again, we omit the index s for brevity) where $L_{-2}, L_{-1}, L_0, L_1, \dots$ are symplectic matrices, and

$$(2.11) \quad L_{-2} = \nu\alpha\alpha^t\sigma, \quad L_{-1} = (\alpha\beta^t + \beta\alpha^t)\sigma \quad (\nu \in \mathbb{C}, \beta \in \mathbb{C}^{2n}).$$

Likewise (2.7) we require

$$(2.12) \quad \beta^t\sigma\alpha = 0.$$

Notice that $\alpha^t\sigma\alpha = 0$ due to the skew-symmetry of the matrix σ .

Further on, we require

$$(2.13) \quad L_0\alpha = k\alpha$$

for some complex number k .

We impose a new relation

$$(2.14) \quad \alpha^t\sigma L_1\alpha = 0.$$

Prove now that the space of Lax operators is closed with respect to the Lie bracket in the case $\mathfrak{g} = \mathfrak{sp}(2n)$ in question. We stress that also in this case there is no structure of associative algebra.

Lemma 2.3. *The properties (2.10)–(2.14) are invariant with respect to the Lie bracket.*

Proof. Let $L'' = [L, L']$.

1) The absence of the terms of orders -4 and -3 in z in L'' follows from the relations $\beta^t \sigma \alpha = 0$, $\alpha^t \sigma \alpha = 0$ (2.12) only.

2) For the order -2 term we have:

$$L''_{-2} = (\nu' k - \nu k' + \beta^t \sigma \beta') \alpha \alpha^t \sigma,$$

hence it has the form required by the relation (2.11) (here and below ν' , β' , L'_i have the same meaning for L' as ν , β , L_i for L).

3) The form of the order -1 term: a straightforward calculation using (2.10), (2.11) gives

$$\begin{aligned} L''_{-1} = & \alpha(\nu \cdot \alpha^t \sigma L'_1 - \nu' \cdot \alpha^t \sigma L_1 + \beta^t \sigma L'_0 - \beta'^t \sigma L_0 + k \beta'^t \sigma - k' \beta^t \sigma) \\ & + (-\nu L'_1 \alpha + \nu' L_1 \alpha - L'_0 \beta + L_0 \beta' + k \beta' - k' \beta) \alpha^t \sigma. \end{aligned}$$

Denote the second bracket by β'' . Then the relations $L_1^t = -\sigma L_1 \sigma^{-1}$, $L_0^t = -\sigma L_0 \sigma^{-1}$ (obeyed by symplectic matrices) imply that the first bracket is equal to $\beta''' \sigma$. Hence

$$L''_{-1} = (\alpha \beta''' + \beta'' \alpha^t) \sigma,$$

where

$$\beta'' = -\nu L'_1 \alpha + \nu' L_1 \alpha - L'_0 \beta + L_0 \beta' + k \beta' - k' \beta.$$

Prove that $\beta''' \sigma \alpha = 0$. Making use of the above expression for $\beta''' \sigma$ we find

$$\beta''' \sigma \alpha = \nu \cdot \alpha^t \sigma L'_1 \alpha - \nu' \cdot \alpha^t \sigma L_1 \alpha + \beta^t \sigma L'_0 \alpha - \beta'^t \sigma L_0 \alpha + k \beta'^t \sigma \alpha - k' \beta^t \sigma \alpha.$$

The first two terms of the just obtained expression vanish by the relation (2.14) applied to L and L' . To the second pair of terms, we apply the relations $L_0 \alpha = k \alpha$ and $L'_0 \alpha = k' \alpha$, after that all remaining terms vanish by the relations (2.12).

4). Prove the relation (2.13) on the eigenvalues of the degree zero term. By definition,

$$(LL')_0 = \nu \alpha \alpha^t \sigma L'_2 + (\alpha \beta^t + \beta \alpha^t) \sigma L'_1 + L_0 L'_0 + L_1 (\alpha \beta'^t + \beta' \alpha^t) \sigma + \nu' L_2 \alpha \alpha^t \sigma.$$

The last pair of terms obviously vanishes having been multiplied by α from the right. We obtain

$$(LL')_0 \alpha = \nu \alpha \alpha^t \sigma L'_2 \alpha + \alpha \beta^t \sigma L'_1 \alpha + \beta \alpha^t \sigma L'_1 \alpha + k k' \alpha.$$

The third summand is equal to 0 by (2.14). Thus α is the eigenvector for a zero degree term even in the product LL' :

$$(LL')_0 \alpha = \alpha(\nu \cdot \alpha^t \sigma L'_2 \alpha + \beta^t \sigma L'_1 \alpha + k k'),$$

and in the commutator we obtain

$$(2.15) \quad L_0''\alpha = \alpha(\nu \cdot \alpha^t \sigma L_2' \alpha - \nu' \cdot \alpha^t \sigma L_2 \alpha + \beta^t \sigma L_1' \alpha - \beta'^t \sigma L_1 \alpha).$$

5). Conservation of the relation $\alpha^t \sigma L_1 \alpha = 0$ in the product and commutator.

For the product, by definition

$$(LL')_0 = L_{-2}L_3' + L_{-1}L_2' + L_0L_1' + L_1L_0' + L_2L_{-1}' + L_3L_{-2}'.$$

Replacing here L_{-2} , L_{-2}' , L_{-1} , L_{-1}' by the known expressions obtain

$$\begin{aligned} \alpha^t \sigma (LL')_0 \alpha &= \nu(\alpha^t \sigma \alpha) \alpha^t \sigma L_3' \alpha + ((\alpha^t \sigma \alpha) \beta^t + (\alpha^t \sigma \beta) \alpha^t) \sigma L_2' \alpha \\ &\quad - k(\alpha^t \sigma L_1' \alpha) + \tilde{k}(\alpha^t \sigma L_1 \alpha) \\ &\quad + \alpha^t \sigma L_2 (\alpha(\beta'^t \sigma \alpha) + \beta'(\alpha^t \sigma \alpha)) + \nu' \alpha^t \sigma L_3 \alpha (\alpha^t \sigma \alpha). \end{aligned}$$

By the relations (2.11)-(2.14) this expression vanishes. $\square$

3. ALMOST GRADED STRUCTURE

In this section we consider the following cases: $\mathfrak{g} = \mathfrak{sl}(n)$, $\mathfrak{g} = \mathfrak{so}(n)$, $\mathfrak{g} = \mathfrak{sp}(2n)$, and, to conclude with, $\mathfrak{g} = \mathfrak{gl}(n)$ which is somewhat special case with respect to the almost gradeness. A general definition of the almost graded structure is given in the introduction. Denote by $\bar{\mathfrak{g}}$ the Lax operator algebra corresponding to $\mathfrak{g}$.

For all above listed algebras $\mathfrak{g}$, except for $\mathfrak{gl}(n)$, and $m \in \mathbb{Z}$ let

$$\mathfrak{g}_m = \{L \in \bar{\mathfrak{g}} \mid (L) + D \geq 0\},$$

where (L) is a divisor of the $\mathfrak{g}$ -valued function L , for $\mathfrak{g} = \mathfrak{sl}(n)$, $\mathfrak{g} = \mathfrak{so}(n)$

$$D = -mP_+ + (m+g)P_- + \sum_{s=1}^{ng} \gamma_s,$$

and for $\mathfrak{g} = \mathfrak{sp}(2n)$

$$D = -mP_+ + (m+g)P_- + 2 \sum_{s=1}^{ng} \gamma_s.$$

We call $\mathfrak{g}_m$ the (homogeneous) subspace of degree m of the Lie algebra $\bar{\mathfrak{g}}$.

Theorem 3.1. For $\mathfrak{g} = \mathfrak{sl}(n)$, $\mathfrak{so}(n)$, $\mathfrak{sp}(2n)$

$$1^\circ. \quad \dim \mathfrak{g}_m = \dim \mathfrak{g}.$$

$$2^\circ. \quad \bar{\mathfrak{g}} = \bigoplus_{m=-\infty}^{\infty} \mathfrak{g}_m.$$

$$3^\circ. \quad [\mathfrak{g}_k, \mathfrak{g}_l] \subseteq \bigoplus_{m=k+l}^{k+l+g} \mathfrak{g}_m.$$

Proof. First prove 1° . For an arbitrary Lie algebra $\mathfrak{g}$, the Riemann-Roch dimension of all $\mathfrak{g}$ -valued functions L satisfying the relation $(L) + D \geq 0$ is equal to $(\dim \mathfrak{g})(ng + 1)$. For $\mathfrak{g} = \mathfrak{sp}(2n)$ it is equal to $(\dim \mathfrak{g})(2ng + 1)$. We will prove that for $\mathfrak{g} = \mathfrak{sl}(n)$, $\mathfrak{g} = \mathfrak{so}(n)$ and an arbitrary $m \in \mathbb{Z}$, there are exactly $\dim \mathfrak{g}$ relations at every pole γ_s , and for $\mathfrak{g} = \mathfrak{sp}(2n)$ the number of those relations is $2 \dim \mathfrak{g}$. This will mean that $\dim \mathfrak{g}_m = \dim \mathfrak{g}$.

Consider first the case $\mathfrak{g} = \mathfrak{sl}(n)$. The elements of the subspace $\mathfrak{g}_m$ satisfy certain conditions of three kinds coming from (2.2), (2.3): on the residues, eigenvalues and traces of the matrix $L \in \mathfrak{g}_m$, as follows:

- 1) at every weak singularity, $L_{-1} = \alpha\beta^t$ which would give $\dim \mathfrak{g}$ relations (since $L_{-1} \in \mathfrak{g}$) if the right hand side was fixed. But the latter depends on the free n -dimensional vector β . Hence we have $\dim \mathfrak{g} - n$ conditions at every one of the ng simple poles γ_s ;
- 2) at every weak singularity $L_0\alpha = k\alpha$ which gives n conditions. Taking account of one free parameter k we obtain $n - 1$ condition at every γ_s ;
- 3) we also have $\text{tr } L = 0$, i.e. one more relation at every weak singularity.

Thus we have $(\dim \mathfrak{g} - n) + (n - 1) + 1 = \dim \mathfrak{g}$ relations at every γ_s , as required.

For $\mathfrak{g} = \mathfrak{so}(n)$ we follow the same line of arguing. Again, the relation $L_{-1} = \alpha\beta^t - \beta\alpha^t$ (2.6) gives $\dim \mathfrak{g} - n$ equations, the relation $L_0\alpha = k\alpha$ (2.8) gives $n - 1$ equation, and $\beta^t\alpha = 0$ (2.7) gives one more equation. Summing up, we obtain $\dim \mathfrak{g}$ equations at every point γ_s .

In the case $\mathfrak{g} = \mathfrak{sp}(2n)$ at every point γ_s we have the following conditions:

$$\begin{aligned} L_{-2} &= \nu\alpha\alpha^t : & \dim \mathfrak{g} - 1 & \text{ conditions (one free parameter } \nu) \\ L_{-1} &= (\alpha\beta^t + \beta\alpha^t)\sigma : & \dim \mathfrak{g} - 2n & \text{ conditions (} 2n \text{ free parameters } \beta) \\ L_0\alpha &= k\alpha : & 2n - 1 & \text{ conditions (one free parameter } k) \\ \beta^t\sigma\alpha &= 0, \alpha^t\sigma L_1\alpha = 0 : & 2 & \text{ conditions,} \end{aligned}$$

i.e. $2 \dim \mathfrak{g}$ conditions at every one of the ng points, as required.

For $m > 0$ and $m < -g$ the subspaces $\mathfrak{g}_m$ are linear independent for the obvious reason: the orders at $P_{\pm}$ of the elements of those subspaces are different for different m .

For $-g \leq m \leq 0$ the linear independence of $\mathfrak{g}_m$ follows from the fact that there are no *everywhere holomorphic* Lax operators. We would like to emphasize that the last argument applies only to the case of simple Lie algebras. That explains why the case of the reductive Lie algebra $\mathfrak{gl}(n)$ requires some modification (see below).

The statement 3° of the theorem follows from the consideration of orders at the points $P_{\pm}$. $\square$

The Theorem 3.1 defines an *almost graded structure* on $\overline{\mathfrak{g}}$.

Consider now the case $\mathfrak{g} = \mathfrak{gl}(n)$. In this case $\overline{\mathfrak{g}}$ contains the subspace of functions taking values in the 1-dimensional space of scalar matrices. Let L be such function. By (2.2) we obtain $\text{tr } L_{-1} = 0$. Since L_{-1} is scalar, we obtain $L_{-1} = 0$. Hence, L is holomorphic except at $P_{\pm}$. Let $\mathcal{A}$ denote the algebra of meromorphic functions on Γ holomorphic except at $P_{\pm}$. Then $L \in \mathcal{A} \cdot id$ where id is the unit matrix. Therefore,

$$(3.1) \quad \overline{\mathfrak{gl}(n)} = \overline{\mathfrak{sl}(n)} \oplus \mathcal{A} \cdot id.$$

In [3], there was introduced a certain base $\{A_m\}$ (later called the *Krichever-Novikov base*) in the space of such functions. Denote $\mathcal{A}_m = \mathbb{C}A_m$ and set $\mathfrak{gl}(n)_m = \mathfrak{sl}(n)_m \oplus (\mathcal{A}_m \cdot id)$. For $m > 0$ and $m < -g$ this definition is equivalent to the above given definition of $\mathfrak{g}_m$ (with $\mathfrak{g} = \mathfrak{gl}(n)$).

With $\mathfrak{g}_m = \mathfrak{gl}(n)_m$, the Theorem 3.1 remains to be true for $\mathfrak{g} = \mathfrak{gl}(n)$, as it follows from (3.1). The relation 3° of the theorem holds with the different upper limit of summation determined by the algebra $\mathcal{A}$ (see [3]).

4. CENTRAL EXTENSIONS OF LAX OPERATOR ALGEBRAS

4.1. Central extensions for the Lax operator algebras over $\mathfrak{gl}(n)$.

The conventional expression for the 2-cocycle giving a central extension for Krichever-Novikov current algebras (in particular, loop algebras) is $\text{tr } \text{res}_{P_{\pm}} L dL'$. For those algebras the cocycle is local. A cocycle χ is called *local* [3] if the constants μ', μ'' exist such that $\chi(\mathfrak{g}_m, \mathfrak{g}_{m'}) = 0$ unless $\mu' \leq m + m' \leq \mu''$ where $\mathfrak{g}_m, \mathfrak{g}_{m'}$ are homogeneous subspaces introduced in the previous section. In case of Lax operator algebras the above cocycle is by no means local. In this section we improve it in order to obtain a local cocycle.

For every operator L the eigenvalue k of its zero degree component L_0 (see (2.3)) can be considered as a linear functional of L . We denote this functional by $k(L)$.

Lemma 4.1. *At every weak singularity, the one-form $\text{tr } L dL'$ has at most simple pole and*

$$(4.1) \quad \text{res } \text{tr } L dL' = k([L, L']).$$

Proof. Calculate both parts of the relation explicitly.

1) Using

$$dL' = -\frac{\alpha\beta''}{z^2} + L'_1 + \dots$$

and (2.1) we obtain

$$L dL' = -\frac{\alpha\beta^t\alpha\beta''}{z^3} - \frac{L_0\alpha\beta''}{z^2} - \frac{L_1\alpha\beta'' - \alpha\beta^t L'_1}{z} + \dots$$

The first term vanishes since $\beta^t\alpha = 0$. The second term vanishes under the symbol of trace, since $L_0\alpha = k\alpha$, and $\text{tr } \alpha\beta'' = \beta''\alpha = 0$. The third term gives us the required residue. We have

$$\text{res } \text{tr } L dL' = \text{tr } (\alpha\beta^t L'_1 - L_1\alpha\beta'') = \beta^t L'_1\alpha - \beta'' L_1\alpha.$$

2) Now calculate the right hand side of the relation in question. Denote by $[L, L']_0$ the zero degree coefficient of the expansion (2.1) for the commutator $[L, L']$. We have

$$\begin{aligned} [L, L']_0 = & \alpha\beta^t L'_1 + L_0 L'_0 + L_1 \alpha\beta'' \\ & - \alpha\beta'' L_1 - L'_0 L_0 - L'_1 \alpha\beta^t. \end{aligned}$$

Multiply both parts of this relation by α from the right hand side. Then the third terms in every row vanish since they contain the factors $\beta''\alpha$, $\beta^t\alpha$ which are zeroes by (2.2). The second terms annihilate since $L_0 L'_0\alpha = k'k\alpha$ and $L'_0 L_0\alpha = kk'\alpha$. Hence

$$[L, L']_0\alpha = \alpha\beta^t L'_1\alpha - \alpha\beta'' L_1\alpha = \alpha(\beta^t L'_1\alpha - \beta'' L_1\alpha).$$

The expression in brackets is a 1×1 -matrix, that is a complex number. Certainly, this complex number is the eigenvalue of $[L, L']_0$ on the vector α , i.e. $k([L, L'])$. Its value exactly meets the above resulting expression for $\text{tr res } L dL'$, which completes the proof. $\square$

We want to eliminate the singularities of the 1-form $\text{tr } LdL'$ at γ_s by subtracting another expression for the eigenvalue on the right hand side of (4.3). Remarkably, we are able to give this new expression in terms of connections in the holomorphic vector bundles on Γ , given in explicit form in [2]. In what follows, $\mathcal{L}$ denotes the 1-form of such connection.

Lemma 4.2. *Let $\mathcal{L}$ be a matrix-valued 1-form such that locally, at a weak singularity,*

$$\mathcal{L} = \mathcal{L}_{-1} \frac{dz}{z} + \mathcal{L}_0 dz + \dots ,$$

and $\mathcal{L}$ satisfies the similar requirements as L (see (2.1)-(2.3)), with the only one modification: $\tilde{\beta}^t \alpha = 1$ where $\mathcal{L}_{-1} = \alpha \tilde{\beta}^t$. Then the 1-form $\text{tr } L\mathcal{L}$ has at most simple pole at $z = 0$ and

$$\text{res tr } L\mathcal{L} = k(L).$$

Proof. We have

$$L = \frac{\alpha \beta^t}{z} + L_0 + \dots , \quad \mathcal{L} = \left(\frac{\alpha \tilde{\beta}^t}{z} + \mathcal{L}_0 + \dots \right) dz,$$

hence

$$L\mathcal{L} = \left(\frac{\alpha \beta^t \alpha \tilde{\beta}^t}{z^2} + \frac{\alpha \beta^t \mathcal{L}_0 + L_0 \alpha \tilde{\beta}^t}{z} + \dots \right) dz.$$

As above, the first term vanishes. For the second term we have

$$\text{res tr } L\mathcal{L} = \text{tr} (\alpha \beta^t \mathcal{L}_0 + L_0 \alpha \tilde{\beta}^t) = \beta^t \mathcal{L}_0 \alpha + \tilde{\beta}^t L_0 \alpha.$$

Further on, $\mathcal{L}_0 \alpha = \tilde{k} \alpha$, $L_0 \alpha = k \alpha$, hence $\beta^t \mathcal{L}_0 \alpha = \tilde{k} \beta^t \alpha = 0$ and $\tilde{\beta}^t L_0 \alpha = k \tilde{\beta}^t \alpha = k$. This completes the proof. $\square$

Theorem 4.3. *For every $\mathcal{L}$ satisfying the conditions of the Lemma 4.2 the 1-form $\text{tr} (L dL' - [L, L'] \mathcal{L})$ is regular except at the points $P_{\pm}$, and the expression*

$$\gamma(L, L') = \text{res}_{P_+} \text{tr} (L dL' - [L, L'] \mathcal{L})$$

gives a local cocycle on the Lax operator algebra.

Proof. In course of proofs of the Lemma 4.1 and the Lemma 4.2, we have seen that the 1-forms $\text{tr } L dL'$ and $\text{tr} [L, L'] \mathcal{L}$ have simple poles at every point γ_s , and their residues are equal to the same quantity $k_s([L, L'])$. Hence, their difference is regular at every γ_s .

Assume, at the point P_+ we have

$$(4.2) \quad L(z) = \sum_{i=m}^{\infty} a_i z^i, \quad L'(z) = \sum_{j=m'}^{\infty} b_j z^j, \quad \mathcal{L}(z) = \sum_{k=m_+}^{\infty} c_k z^k dz.$$

Then

$$L(z) dL'(z) = \sum_{p=m+m'}^{\infty} \left(\sum_{i+j=p} j a_i b_j \right) z^{p-1} dz,$$

and

$$[L(z), L'(z)]\mathcal{L} = \sum_{p=m+m'+m_+}^{\infty} \left(\sum_{i+j+k=p} [a_i, b_j]c_k \right) z^p dz.$$

In order one of these 1-forms had a nontrivial residue at the point P_+ it is necessary that either $m + m' \leq 0$ or $m + m' + m_+ \leq -1$, in other words

$$m + m' \leq \max\{0, -1 - m_+\}.$$

If L and L' are homogeneous of degrees m, m' then at the point P_- their expansions (similar to (4.2)) begin with $i = -m - g, j = -m' - g$, respectively. The expansion for $\mathcal{L}$ begins with some integer m_- . Hence the condition at P_- looks as follows:

$$-m - m' - 2g \leq \max\{0, -1 + m_-\}.$$

Finally, we obtain

$$\min\{0, 1 - m_-\} - 2g \leq m + m' \leq \max\{0, -1 - m_+\}.$$

Since $m_{\pm}$ are fixed ($\mathcal{L}$ is fixed), the latter exactly means that the cocycle is local. $\square$

4.2. Central extensions for the Lax operator algebras over $\mathfrak{so}(n)$. We keep here the same line of arguing as in the previous section.

Lemma 4.4. *At every weak singularity, the one-form $\text{tr } L dL'$ has at most simple pole and*

$$(4.3) \quad \text{res } \text{tr } L dL' = 2k([L, L']).$$

Proof. 1) Using (2.5) and the relation

$$dL' = -L'_{-1}z^{-2} + L'_1 + \dots,$$

where L'_{-1} is given by (2.6), we obtain

$$(4.4) \quad L dL' = -\frac{L_{-1}L'_{-1}}{z^3} - \frac{L_0L'_{-1}}{z^2} - \frac{L_1L'_{-1} - L_{-1}L'_1}{z} + \dots.$$

For the first term we have

$$L_{-1}L'_{-1} = (\alpha\beta^t - \beta\alpha^t)(\alpha\beta'^t - \beta'\alpha^t) = \alpha(\beta^t\alpha)\beta'^t - \beta(\alpha^t\alpha)\beta'^t - \alpha\beta^t\beta'\alpha^t - \beta\alpha^t\beta'\alpha^t.$$

The first two summands vanish due to (2.7). For the remainder, we have

$$\text{tr}(L_{-1}L'_{-1}) = \text{tr}(-\alpha\beta^t\beta'\alpha^t - \beta\alpha^t\beta'\alpha^t) = -(\alpha^t\alpha)\beta^t\beta' - (\alpha^t\beta)\alpha^t\beta'$$

which again vanishes due to (2.7).

Again (as in Section 4.1) the term containing z^{-2} vanishes under the symbol of trace. By definition,

$$L_0L'_{-1} = L_0(\alpha\beta'^t - \beta'\alpha^t) = k\alpha\beta'^t - L_0\beta'\alpha^t.$$

Now observe that $\text{tr}(L_0\beta'\alpha^t) = \text{tr}(\alpha^t L_0\beta')$ and $\alpha^t L_0 = -k\alpha^t$. Hence $\text{tr}(L_0 L'_{-1}) = 2k\alpha^t\beta'$ which vanishes due to (2.7).

The third term in (4.4) gives us the required residue. We have

$$\text{res } L dL' = (L_{-1}L'_1 - L_1L'_{-1}).$$

Substituting L_{-1} , L'_{-1} from (2.6), we have

$$\text{res } L dL' = \alpha\beta^t L'_1 - \beta\alpha^t L'_1 - L_1\alpha\beta'^t + L_1\beta'\alpha^t$$

hence

$$\text{tr res } L dL' = \beta^t L'_1 \alpha - \alpha^t L'_1 \beta - \beta'^t L_1 \alpha + \alpha^t L_1 \beta'.$$

Due to the skew-symmetry of matrices L_1 , L'_1 the first two summands of the last relation are equal, and the same about the last two summands. Hence

$$\text{tr res } L dL' = 2(\beta^t L'_1 \alpha - \beta'^t L_1 \alpha).$$

From (2.9) obviously follows that $[L, L']_0 \alpha = \beta^t L'_1 \alpha - \beta'^t L_1 \alpha$ which proves the lemma. $\square$

Lemma 4.5. *Let $\mathcal{L}$ be a skew-symmetric matrix-valued 1-form such that locally, at a weak singularity,*

$$\mathcal{L} = \mathcal{L}_{-1} \frac{dz}{z} + \mathcal{L}_0 dz + \dots$$

where $\mathcal{L}_{-1} = \alpha\tilde{\beta}^t - \tilde{\beta}\alpha^t$, $\tilde{\beta}^t\alpha = 1$ and $\mathcal{L}_0\alpha = \tilde{k}\alpha$. Then the 1-form $\text{tr } L\mathcal{L}$ has at most simple pole at $z = 0$ and

$$\text{res tr } L\mathcal{L} = 2k(L).$$

Proof. The coefficient of the degree -2 term in the product $L\mathcal{L}$ is equal, by definition, to

$$(\alpha\beta^t - \beta\alpha^t)(\alpha\tilde{\beta}^t - \tilde{\beta}\alpha^t) = \alpha(\beta^t\alpha)\tilde{\beta}^t - \alpha\beta^t\tilde{\beta}\alpha^t - \beta(\alpha^t\alpha)\tilde{\beta}^t + \beta\alpha^t\tilde{\beta}\alpha^t.$$

The first and the third terms vanish due to (2.7). For the trace of the remainder we have

$$\text{tr}(-\alpha\beta^t\tilde{\beta}\alpha^t + \beta\alpha^t\tilde{\beta}\alpha^t) = -(\alpha^t\alpha)\beta^t\tilde{\beta} + (\alpha^t\beta)\alpha^t\tilde{\beta}$$

which vanishes for the same reason.

Multiplying expansions for L and $\mathcal{L}$ we find

$$\text{tr res}(L\mathcal{L}) = \text{tr}(L_{-1}\mathcal{L}_0 + L_0\mathcal{L}_{-1}) = \text{tr}(\alpha\beta^t - \beta\alpha^t)\mathcal{L}_0 + \text{tr } L_0(\alpha\tilde{\beta}^t - \tilde{\beta}\alpha^t).$$

For the first summand we have

$$\text{tr}(\alpha\beta^t - \beta\alpha^t)\mathcal{L}_0 = \text{tr}(\alpha\beta^t\mathcal{L}_0 - \beta\alpha^t\mathcal{L}_0) = \beta^t\mathcal{L}_0\alpha - \alpha^t\mathcal{L}_0\beta.$$

Observe that, by skew-symmetry, $\alpha^t \mathcal{L}_0 \beta = -\beta^t \mathcal{L}_0 \alpha$, hence

$$\mathrm{tr}(\alpha \beta^t - \beta \alpha^t) \mathcal{L}_0 = \beta^t \mathcal{L}_0 \alpha - \alpha^t \mathcal{L}_0 \beta = 2\beta^t \mathcal{L}_0 \alpha = 2\tilde{k} \beta^t \alpha$$

which vanishes due to (2.7).

For the second summand we have

$$\mathrm{tr} L_0(\alpha \tilde{\beta}^t - \tilde{\beta} \alpha^t) = \mathrm{tr}(L_0 \alpha \tilde{\beta}^t - L_0 \tilde{\beta} \alpha^t) = \tilde{\beta}^t L_0 \alpha - \alpha^t L_0 \tilde{\beta}.$$

Since $L_0 \alpha = k\alpha$, $\alpha^t L_0 = -k\alpha^t$ and $\tilde{\beta}^t \alpha = \alpha^t \tilde{\beta} = 1$ we obtain

$$\mathrm{tr} L_0(\alpha \tilde{\beta}^t - \tilde{\beta} \alpha^t) = 2k.$$

□

Theorem 4.6. *For every $\mathcal{L}$ satisfying the conditions of the Lemma 4.5 the 1-form $\mathrm{tr}(L dL' - [L, L'] \mathcal{L})$ is regular except at the points $P_{\pm}$, and the expression*

$$\gamma(L, L') = \mathrm{res}_{P_+} \mathrm{tr}(L dL' - [L, L'] \mathcal{L})$$

gives a local cocycle on the Lax operator algebra.

The proof is quite similar to the proof of the Theorem 4.3. It relies only on the absence of residues of the 1-form defining the cocycle at the weak singularities.

There is a certain ambiguity in the definition of $\mathcal{L}$ in Lemma 4.5. For example, we could require that $\mathcal{L}_{-1} = \alpha \tilde{\beta}^t$ and take $L dL' - 2[L, L'] \mathcal{L}$ in the Theorem 4.6.

4.3. Central extensions for the Lax operator algebras over $\mathfrak{sp}(2n)$.

We keep the same line of arguing as in the previous sections. First, prove the following analogue of the Lemma 4.4.

Lemma 4.7. *At every weak singularity the 1-form $\mathrm{tr} L dL'$ has at most simple pole, and*

$$(4.5) \quad \mathrm{res} \mathrm{tr} L dL' = 2k([L, L']).$$

Proof. A straightforward calculation of $L dL'$ based on the expansion (2.10) shows that the terms with z^{-5} and z^{-4} of that matrix-valued 1-form are equal to 0 by the relations (2.12).

For the term with z^{-3} we have

$$(L dL')_{-3} = -(\beta^t \sigma \beta' - 2\nu') \alpha \alpha^t \sigma.$$

This expression vanishes under the symbol of trace:

$$\mathrm{tr}(L dL')_{-3} = -(\beta^t \sigma \beta' - 2\nu')(\alpha^t \sigma \alpha) = 0.$$

Similarly, using (2.12)-(2.14) we have

$$\mathrm{tr} (LdL')_{-2} = \nu(\alpha^t \sigma L'_1 \alpha) - 2k(\beta'^t \sigma \alpha) - \nu'(\alpha^t \sigma L_1 \alpha) = 0.$$

Thus, the 1-form $\mathrm{tr} LdL'$ indeed has at most simple pole at the point in question. A straightforward calculation results in

$$\mathrm{tr} (LdL')_{-1} = 2(\nu \cdot \alpha^t \sigma L'_2 \alpha - \nu' \cdot \alpha^t \sigma L_2 \alpha + \beta^t \sigma L'_1 \alpha - \beta'^t \sigma L_1 \alpha)$$

which exactly coincides with the double expression (2.15) for $k([L, L'])$. $\square$

Lemma 4.8. *Let $\mathcal{L}$ be a $\mathfrak{g}$ -valued 1-form such that locally, in the neighborhood of any weak singularity,*

$$\mathcal{L} = \mathcal{L}_{-1} \frac{dz}{z} + \mathcal{L}_0 dz + \dots$$

where $\mathcal{L}_{-1} = (\alpha \tilde{\beta}^t + \tilde{\beta} \alpha^t) \sigma$, $\tilde{\beta}^t \sigma \alpha = 1$, $\mathcal{L}_0 \alpha = \tilde{k} \alpha$, $\alpha^t \sigma \mathcal{L}_1 \alpha = 0$. Then the 1-form $\mathrm{tr} L\mathcal{L}$ has at most simple pole at $z = 0$

$$\mathrm{res} \mathrm{tr} L\mathcal{L} = 2k(L).$$

Proof. The expansion for $L\mathcal{L}$ begins with z^{-3} . We have

$$\mathrm{tr} (L\mathcal{L})_{-3} = \nu(\tilde{\beta}^t \sigma \alpha)(\alpha^t \sigma \alpha) = 0,$$

$$\mathrm{tr} (L\mathcal{L})_{-2} = \nu(\alpha^t \sigma \tilde{\beta})(\alpha^t \sigma \alpha) + (\beta^t \sigma \tilde{\beta})(\alpha^t \sigma \alpha) + (\alpha^t \sigma \tilde{\beta})(\alpha^t \sigma \beta) = 0.$$

Thus, the 1-form $\mathrm{tr} L\mathcal{L}$ indeed has at most simple pole at the point in question. The calculation of the residue results in

$$\mathrm{tr} (L\mathcal{L})_{-1} = \nu \cdot \alpha^t \sigma \mathcal{L}_1 \alpha + \beta^t \sigma (\mathcal{L}_0 \alpha) + (\alpha^t \sigma \mathcal{L}_0) \beta + \tilde{\beta}^t \sigma (L_0 \alpha) + (\alpha^t \sigma L_0) \tilde{\beta}.$$

By assumption of the lemma, $\alpha^t \sigma \mathcal{L}_1 \alpha = 0$, $\mathcal{L}_0 \alpha = \tilde{k} \alpha$. The last implies also that $\alpha^t \sigma \mathcal{L}_0 = -\tilde{k} \alpha^t \sigma$. For this reason

$$\mathrm{tr} (L\mathcal{L})_{-1} = \tilde{\beta}^t \sigma (L_0 \alpha) + (\alpha^t \sigma L_0) \tilde{\beta}.$$

By the relations $L_0 \alpha = k \alpha$, $\alpha^t \sigma L_0 = -k \alpha^t \sigma$ we have

$$\mathrm{tr} (L\mathcal{L})_{-1} = 2k(\tilde{\beta}^t \sigma \alpha) = 2k.$$

$\square$

As above, the last two lemmas imply

Theorem 4.9. *For every $\mathcal{L}$ satisfying the conditions of the Lemma 4.8 the 1-form $\mathrm{tr} (LdL' - [L, L']\mathcal{L})$ is regular except at the points $P_{\pm}$, and the expression*

$$\gamma(L, L') = \mathrm{res}_{P_{+}} \mathrm{tr} (LdL' - [L, L']\mathcal{L})$$

gives a local cocycle on the Lax operator algebra.

REFERENCES

- [1] I.M.Krichever *Vector bundles and Lax equations on algebraic curves*. Commun.Math.Phys. 229 (2002), p.229-269. hep-th/0108110.
- [2] I.M.Krichever *Isomonodromy equations on algebraic curves, canonical transformations and Witham equations*. hep-th/0112096.
- [3] Krichever, I.M, Novikov, S.P., *Algebras of Virasoro type, Riemann surfaces and structures of the theory of solitons*. Funktional Anal. i. Prilozhen. **21**, No.2 (1987), 46-63.
- [4] Krichever, I.M, Novikov, S.P., *Virasoro type algebras, Riemann surfaces and strings in Minkowski space*. Funktional Anal. i. Prilozhen. **21**, No.4 (1987), 47-61.
- [5] Krichever, I.M, Novikov, S.P., *Algebras of Virasoro type, energy-momentum tensors and decompositions of operators on Riemann surfaces*. Funktional Anal. i. Prilozhen. **23**, No.1 (1989), 46-63.
- [6] Krichever, I.M, Novikov S.P., *Holomorphic bundles and commuting difference operators. Two-point constructions*. Usp. Mat. Nauk, **55**, No.4 (2000) 181-182.
- [7] Krichever I.M., Novikov S.P. *Holomorphic bundles on Riemann surfaces and Kadomtsev-Petviashvili equation (KP). I*. Funkt. Analys i Pril. , **12** (1978), 4, 41-52.
- [8] Krichever I.M., Novikov S.P. *Holomorphic bundles on algebraic curves and nonlinear equations*. Uspekhi Math. Nauk (Russ. Math.Surv), **35** (1980), 6, 47-68.
- [9] Schlichenmaier, M., *Local cocycles and central extensions for multi-point algebras of Krichever-Novikov type*. J. Reine und Angewandte Mathematik **559**(2003), 53-94.
- [10] Schlichenmaier, M., *Higher genus affine algebras of Krichever-Novikov type*. Moscow Math. J. **3** (4) (2003), 1395-1427, math/0210360.
- [11] Sheinman, O.K., *Affine Krichever-Novikov algebras, their representations and applications*. In: Geometry, Topology and Mathematical Physics. S.P.Novikov's Seminar 2002-2003, V.M.Buchstaber, I.M.Krichever, eds. AMS Translations (2) 212, p.p. 297-316. math.RT/0304020
- [12] Tyurin, A.N. *Classification of vector bundles on an algebraic curve of an arbitrary genus*. Soviet Izvestia, ser. Math., **29** (1965), 657-688.

(Igor M. Krichever) LANDAU INSTITUTE FOR THEORETICAL PHYSICS, UL.KOSYGINA 2, MOSCOW, 119334, RUSSIA, AND COLUMBIA UNIVERSITY, 2960 BROADWAY, NEW YORK, NY 10027-6902, USA
E-mail address: krichev@math.columbia.edu

(Oleg K. Sheinman) STEKLOV MATHEMATICAL INSTITUTE, UL. GUBKINA, 8, MOSCOW, 119991 RUSSIA AND INDEPENDENT UNIVERSITY OF MOSCOW, BOLSHOI VLASIEVSKII PER. 11, MOSCOW, RUSSIA
E-mail address: sheinman@mi.ras.ru